

SANTE MENTALE DES MIGRANTS

A. TORTELLI

Praticien hospitalier, CAPSYS, Pôle Psychiatrie Précarité GHU - Paris
INSERM IPLESP
Institut Convergences Migrations, département Health

2024




Constats


Depuis le 19ème siècle, à travers les différentes vagues migratoires on observe un risque accru de troubles psychiatriques par rapport aux natifs

Ce risque est augmenté aussi chez les descendants


Néanmoins, le risque varie en fonction des origines, pays d'accueil, type de migration, conditions de vie, etc



Comment expliquer?



Etiologie des troubles psychiatriques: Interactions gène-environnement



Gandal et al, 2018

Santé et immigration: 12 Juin 2019

L' expérience migratoire
augmente l'exposition à
des facteurs de risque
environnementaux (bio-
psycho-sociaux) des
troubles psychiatriques



Etudes en Population générale

Troubles psychiatriques : cohorte en population danoise, migrants et descendants (Cantor Graae & Pedersen, 2013)

Troubles psychotiques

Incidence :

McGrath 2004, 55 études, 33 pays (statut migratoire OR= 4.6)

Bourque, 2011 (2x), 1^{ère} et deuxième génération, différence ethnique

Prévalence: Saha et al (2005) : 188 études, 46 pays OR = 1.8

Troubles de l'humeur

Swinnen & Selten, 2018: OR=1.38 (CI 1.17–1.62)

Demandeurs d'asyle/refugiés

Prévalence:

Metanalyse 1 : (Blackmore et al, 2020)

PTSD : 31.46%

Depression 31.5%

Tb Anxieux 11%

Psychose 1.51%

Metanalyse 2 : (Fazel et al, 2005)

PTSD 9%

Depression 5%

Psychose 2%

Prevalence of serious mental disorder in 7000 refugees resettled in western countries: a systematic review

Mina Fazel, Jeremy Wheeler, John Danesh

Summary

Background About 13 million people are classified as refugees worldwide, and many more former refugees have been granted citizenship in their new countries. However, the prevalence of post-traumatic stress disorder, major depression, or psychotic illnesses in these individuals is not known. We did a systematic review of surveys about these disorders in general refugee populations in western countries.

Methods We searched for psychiatric surveys that were based on interviews of unselected refugee populations and that included current diagnoses of post-traumatic stress disorder, major depression, psychotic illnesses, or generalised anxiety disorder. We did computer-assisted searches, scanned reference lists, searched journals, and corresponded with authors to determine prevalence rates of these mental disorders and to explore potential sources of heterogeneity such as diagnostic criteria, sampling methods, and other characteristics.

RESEARCH ARTICLE

The prevalence of mental illness in refugees and asylum seekers: A systematic review and meta-analysis

Rebecca Blackmore^①, Jacqueline A. Boyle^②, Mina Fazel^③, Sanjeeva Ranasingha^④, Kylie M. Gray^{③,4}, Grace Fitzgerald^①, Marie Misso^①, Melanie Gibson-Helm^{①*}

1 Monash Centre for Health Research and Implementation, School of Public Health and Preventive Medicine, Monash University, Melbourne, Australia, **2** Department of Psychiatry, Oxford University, Oxford, United Kingdom, **3** Centre for Developmental Psychiatry & Psychology, Department of Psychiatry, School of Clinical Sciences, Monash University, Melbourne, Australia, **4** Centre for Educational Development, Appraisal and Research (CEDAR), University of Warwick, Coventry, United Kingdom

* melanie.gibson@monash.edu

En France

Population générale

Troubles psychotiques

Incidence: statut migratoire: OR=1.4 et descendants, différence ethnique (Tortelli et al, 2021)

Prévalence: 3 générations (Amad et al, 2013)

Troubles dépressifs : OR=1.4 (Pignon et al, 2017)

En France

Populations spécifiques

SAMENTA (sans abri): Dépression : 26% (3x), PTSD : 6% (6x), Psychose associée à un séjour > 10 ans en France (Laporte et al, 2018; Tortelli et al, 2022)

ENFAMS (hébergement) : 29% dépression, 17% PTSD (12 derniers mois) (Vandendorren et al, 2016)

Premiers Pas (AME): 1 sur 6 = PTSD, association entre précarité et PTSD (Prieur et al, 2022)

Vague actuelle: facteurs additionnels de vulnérabilité

- Migration forcée
- Trajet (Lybie, Bateau)
- Langue
- MNA
- Familles
- Politiques migratoires



CLINIQUE des exilés : superposition de symptômes

Trauma  PTSD

- Tb du sommeil, cauchemars
- Ruminations anxieuses, attaques de panique
- Idées suicidaires
- Hallucinations
- Bégayements, crises dissociatives,
« somatisations »

CLINIQUE: évolution

Évolution en fonction de facteurs internes et externes :
en dents de scie, avec des moments de crise
(notamment des idées suicidaires)

Facteurs externes d'impact négatif:


Lenteur de la procédure

Inactivité

Isolement

Mobilité forcée

CLINIQUE: Prise en charge



Des situations complexes :
prise en charge globale, avec identification des priorités (médico-sociales) avec la personne



Traitements: bonne réponse aux traitements pharmacologiques et aux psychothérapies de type TCC, EMDR, etc



pas de savoir spécifique



Impact des activités sportives, de méditation, de socialisation et d'acculturation

REFERENCES

- Bryant, R. A., Schafer, A., Dawson, K. S., Anjuri, D., Mulili, C., Ndogoni, L., Koyiet, P., Sijbrandij, M., Ulate, J., Harper Shehadeh, M., Hadzi-Pavlovic, D. and van Ommeren, M. (2017) 'Effectiveness of a brief behavioural intervention on psychological distress among women with a history of gender-based violence in urban Kenya: A randomised clinical trial', *PLoS Medicine*, 14(8), e1002371.
- Burnett A. P. M. (2001) 'Health needs of asylum seekers and refugees. :544-7', *British Medical Journal*, 322, 544-547.
- Dawson, K. S., Bryant, R. A., Harper, M., Kuowei Tay, A., Rahman, A., Schafer, A. and van Ommeren, M. (2015) 'Problem Management Plus (PM+): a WHO transdiagnostic psychological intervention for common mental health problems', *World Psychiatry*, 14(3), 354-357.
- Foa, E. B. (1995) 'Posttraumatic stress diagnostic scale (PDS)', *Minneapolis: National Computer Systems*.
- Guisao, A., Quéré, M. and Riou, F. (2017) Profil socio-Sanitaire et besoins de suivi sanitaire des personnes Accueillies dans les structures d'Île-de-France hébergeant des migrants évacués de campements de fortune parisiens, Observatoire du Samu social.
- Hamdani, S. U., Ahmed, Z., Sijbrandij, M., Nazir, H., Masood, A., Akhtar, P., Amin, H., Bryant, R. A., Dawson, K., van Ommeren, M., Rahman, A. and Minhas, F. A. (2017) 'Problem Management Plus (PM+) in the management of common mental disorders in a specialized mental healthcare facility in Pakistan; study protocol for a randomized controlled trial', *Int J Ment Health Syst*, 11, 40.
- Johnston, V. (2009) 'Australian asylum policies: have they violated the right to health of asylum seekers?', *Australian and New Zealand Journal of Public Health*, 33(1), 40-46.
- Lamkadem, M., Stronks, K., Devillé, W. D., Olff, M., Gerritsen, A. A. and Essink-Bot, M.-L. (2014) 'Course of post-traumatic stress disorder and health care utilisation among resettled refugees in the Netherlands', *BMC Psychiatry*, 14(1), 90.
- Nickerson, A., Bryant, R. A., Silove, D. and Steel, Z. (2011) 'A critical review of psychological treatments of posttraumatic stress disorder in refugees', *Clinical psychology review*, 31(3), 399-417.
- Nosè, M., Ballette, F., Bighelli, I., Turrini, G., Purgato, M., Tol, W., Priebe, S. and Barbui, C. (2017) 'Psychosocial interventions for post-traumatic stress disorder in refugees and asylum seekers resettled in high-income countries: Systematic review and meta-analysis', *PLoS One*, 12(2), e0171030.
- Siffert, I., Cordone, A., Réginal, M. and Le Méner, E. (2018) *L'accès aux soins des « migrants » en Île-de-France* Observatoire du Samu social.
- Silove, D., Sinnerbrink, I., Field, A., Manicavasagar, V. and Steel, Z. (1997) 'Anxiety, depression and PTSD in asylum-seekers: associations with pre-migration trauma and post-migration stressors', *The British Journal of Psychiatry*, 170(4), 351-357.
- Toar, M., O'Brien, K. K. and Fahey, T. (2009) 'Comparison of self-reported health & healthcare utilisation between asylum seekers and refugees: an observational study', *BMC Public Health*, 9:214., 214.
- Tortelli, A., Perquier, F., Melchior, M., Lair, F., Encatassamy, F., Masson, C., ... & Mercuel, A. (2020). Mental Health and Service Use of Migrants in Contact with the Public Psychiatry System in Paris. *International Journal of Environment*
- Vandendorren, S., Le Mener, E., Oppenchaïm, N., Arnaud, A., Jangal, C., Caum, C., ... & Guyavarch, E. (2016). Characteristics and health of homeless families: the ENFAMS survey in the Paris region, France 2013. *The European Journal of Public Health*, 26(1), 71-76.al Research and Public Health, 17(24), 9397.

